

Referral Status: New Patient Updated Patient Information

CMC Infusion Center- Referral Face Sheet



Patient Name: _____ Patient D.O.B. _____

Current Height: _____ Current Weight _____

Patient Phone # (_____) _____ - _____ Gender _____

Patient Address _____

City _____ State _____ Zip Code _____

Related Diagnosis Code (ICD-10 code): _____

Referring Provider (Please Print) _____

Office Phone # (_____) _____ - _____ Office Contact: _____

If referring provider and/or patient is external to Conway Medical Center. Please include:

- Most recent history and physical including:
 - Comprehensive medication list
 - Past medical history
 - Related past/ failed therapies (with dates)
 - Known allergies
- Most recent labs / test result (Ex. CBC, PPD, Hepatitis screening, etc.)
- Insurance Information

Please fax requested documentation, face sheet, and treatment plan to **843-234-5460** attention CMC Medication Management Specialist. Please feel free to call 843-234-8575 with any questions.

We look forward to the opportunity to be a part of your patient's treatment journey at Conway Medical Center.

Referral Status New Order Change Order Renewal



Infliximab Treatment Plan (Rheumatology & Neurology)

Patient Name: _____ Patient DOB _____

General

- Diagnosis- Rheumatoid Arthritis with Rheumatoid factor ICD-10 Code _____
- Diagnosis- Rheumatoid Arthritis without Rheumatoid factor ICD-10 Code _____
- Diagnosis- Ankylosing Spondylitis ICD-10 Code _____
- Diagnosis- Sarcoidosis of the lung ICD-10 Code _____
- Diagnosis- Psoriatic Arthropathy ICD-10 Code _____
- Diagnosis- Other _____ ICD-10 Code _____

Pt. Weight _____ Pt Height _____ Known Allergies _____

Requested Start Date ____/____/____

Has patient previously received this medication No Yes, if so, date of last infusion ____/____/____

Authorized Treatment Duration 12 Months Other _____

***Note: Pharmacy will interchange biosimilar product per insurance approval unless box on page 4 is selected prohibiting interchange. If interchange is **NOT** allowed by provider, please indicate preferred product here:
Preferred Product: _____

Pharmacy may round dose to nearest 100mg

Infuse over 2 hours for first 3 infusions. If / when patient tolerates at least three consecutive infusions over 2 hours without signs or symptoms of intolerance, hypersensitivity, or anaphylactic reaction, nursing may request pharmacy to adjust rate to run over 1 hour for subsequent infusions

Infusion (Month 1 to 12) Duration: 12 Months

Vital Signs (Month 1 to 12)
every 30 min, Resp, BP, HR, Temperature on admission to the unit and prior to infusion HR, BP every 30 minutes during the infusion and prior to discharge

Pre- Treatment Medications

Tylenol (Month 1 to 12)

650 mg, Oral, Tab, Day of Tx
Comments: Administer 30 minutes before infliximab dose

Prescriber Signature (No Stamped Signatures or Electronic Signatures)

Provider Signature _____ Provider Name (please print) _____

Date ____/____/____ Patient Name _____ Patient DOB _____

Infliximab Treatment Plan (Rheumatology & Neurology)

Benadryl (Month 1 to 12)

- 25 mg, IV Push, Injection, Day of Tx
Comments: Administer 30 minutes before infliximab dose
- 50 mg, IV Push, Injection, Day of Tx
Comments: Administer 30 minutes before infliximab dose

SOLU-Medrol (Month 1 to 12)

- 40 mg, IV Push, Powder-Inj, Day of Tx
Comments: Administer 30 minutes before infliximab dose
- 125 mg, IV Push, Powder-Inj, Day of Tx
Comments: Administer 30 minutes before infliximab dose

Loading Dose Infliximab (week 0,2,6)

- 3 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 4 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 5 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 6 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 7 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 8 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 9 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 10 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less

Prescriber Signature (No Stamped Signatures or Electronic Signatures)

Provider Signature _____ Provider Name (please print) _____

Date ____ / ____ / ____ Patient Name _____ Patient DOB _____

Infliximab Treatment Plan (Rheumatology & Neurology)

Maintenance Dose Infliximab

Every 8 weeks Every 6 weeks Every 4 weeks Other -Every ____ Weeks

- 3 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 4 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 5 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 6 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 7 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 8 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 9 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less
- 10 mg/kg, IV Piggyback, Injection, Day of Tx
Comments: in 250 mL NS. Administer IV over at least 2 hours with an in-line, low-protein-binding filter 1.2 µm or less

Communication Order (Month 1 to 12)
Patient must stay in infusion area for 30 min post infusion. If vital signs normal may be discharged to home. If abnormal, call physician. Patient should be accompanied by another adult upon discharge

Communication Order (Month 1 to 12)
Ensure results of Hepatitis B surface antigen testing and record results if not present in medical record. Results must be within last 12 months.

Communication Order (Month 1 to 12)
Ensure negative PPD or other test to exclude latent tuberculosis. Results must be within the last 12 months.

Notify Provider (Month 1 to 12)
*for fever/chills, chest pain, hypotension, hypertension, dyspnea, Pruritis, urticaria, persistent flushing
Temperature > 100 or HR <50 or > 130*

Prescriber Signature (No Stamped Signatures or Electronic Signatures)

Provider Signature _____ Provider Name (please print) _____

Date ____ / ____ / ____ Patient Name _____ Patient DOB _____

Infliximab Treatment Plan (Rheumatology & Neurology)

- Notify Provider (Month 1 to 12)
For POSITIVE results of PPD test or other test to exclude latent tuberculosis. Results must be within the last 12 months.
- Notify Provider (Month 1 to 12)
For POSITIVE results of Hepatitis B surface antigen or if results are not available. Results must be within the last 12 months.
- INF Infusion Room Orders Subphase (Month 1 to 12)

Labs (Month 1 to 12) Duration: 12 Months

- Quantiferon(R)-TB Gold, 4T, Incu-(36971) (If Needed)
- Hepatitis B Surface Antibody, Qual-(499) (If Needed)
- Hepatitis B Surface Antigen w Ref/Conf-(498) (If Needed)
- Hepatitis B Core Antibody, Total-(501) (If Needed)
- CBC w/ Diff (If Needed)
- Sedimentation Rate (If Needed)
- C-Reactive Protein (If Needed)
- BUN (If Needed)
- ALT(If Needed)
- AST (If Needed)
- BMP (If Needed)
- Urinalysis with Microscopic, if indicated (If Needed)

Documentation of medications attempted and failed:

- Steroids NSAIDS

Methotrexate Documentation

- Used in combination with infliximab Patient Intolerant Patient has contraindication

Disease Modifying/ Antirheumatic Drugs

- Arava Plaquenil Alzulfidine Other _____

Biologics

- Orencia Humira Cimzia Enbrel Simponi Actemra Xeljanz Other _____

- Biosimilar Substitution **NOT** permitted

Prescriber Signature (No Stamped Signatures or Electronic Signatures)

Provider Signature _____ Provider Name (please print) _____

Date ____ / ____ / ____ Patient Name _____ Patient DOB _____